

MATH 618 (SPRING 2025, PHILLIPS): HOMEWORK 5

This assignment is due on Canvas on Monday 5 May 2025 at 9:00 pm.

Problem 1 (Problem 4 in Chapter 10 of Rudin's book). Let f be an entire function. Suppose that there are constants $A, B > 0$ and $k \in \mathbb{Z}_{>0}$ such that $|f(z)| \leq A + B|z|^k$ for all $z \in \mathbb{C}$. Prove that f is a polynomial.

Problem 2 (Problem 6 in Chapter 10 of Rudin's book). Prove that there is a region Ω such that $\exp(\Omega) = B_1(1)$. Prove that there are many such choices of Ω . Prove that, for any such choice of Ω , the restriction $\exp|_{\Omega}$ is injective. Fix one such choice of Ω , and define $\log: B_1(1) \rightarrow \Omega$ to be the inverse function of $\exp|_{\Omega}$. Prove that $\log'(z) = z^{-1}$. Find the coefficients a_n in the expansion

$$\frac{1}{z} = \sum_{n=0}^{\infty} a_n (z-1)^n,$$

and hence find the coefficients c_n in the expansion

$$\log(z) = \sum_{n=0}^{\infty} c_n (z-1)^n.$$

In which other disks can this be done?

You may use the standard facts about $\exp$, $\log$, $\sin$, and $\cos$ in calculus of a real variable, the formula $\exp(x + iy) = e^x(\cos(y) + i \sin(y))$ for $x, y \in \mathbb{R}$, and the polar form $z = r \exp(i\theta)$ of a complex number z , with unique $r \geq 0$ and, if $r > 0$, uniquely determined $\theta \bmod 2\pi\mathbb{Z}$. You may also use the following facts about $\exp$:

- (1) $\exp(z_1 + z_2) = \exp(z_1)\exp(z_2)$ for every $z_1, z_2 \in \mathbb{C}$. (This can be gotten from the power series.)
- (2) $\exp$ is periodic with period $2\pi i$. (This follows from the formula $\exp(x + iy) = e^x(\cos(y) + i \sin(y))$.)
- (3) For every $z \in \mathbb{C} \setminus \{0\}$, there is $b \in \mathbb{C}$ such that $\exp(b) = z$. (Write $z = r \exp(i\theta)$ with $r \in (0, \infty)$ and $\theta \in \mathbb{R}$. Then take $b = \log(r) + i\theta$, using the usual definition of $\log: (0, \infty) \rightarrow \mathbb{R}$.)

To keep the amount of writing down, I suggest writing a unified proof for all possible choices of the disk.

The following problem counts as two ordinary problems.

Problem 3 (Problem 17 in Chapter 10 of Rudin's book). Determine the largest regions in which the following functions are defined and holomorphic:

$$(1) \quad f(z) = \int_0^1 \frac{1}{1+tz} dt.$$

$$(2) \quad g(z) = \int_0^{\infty} \frac{e^{tz}}{1+t^2} dt.$$

Date: 5 May 2025.

$$(3) \quad h(z) = \int_{-1}^1 \frac{e^{tz}}{1+t^2} dt.$$

Hint: Use Problem 16 in Chapter 10 of Rudin's book (in the previous homework assignment), or combine Morera's Theorem and Fubini's Theorem. In either case, be sure to verify the hypotheses of the theorems you use.

Problem 4 (Problem 3 in Chapter 10 of Rudin's book). Suppose that f and g are entire functions, and that $|f(z)| \leq |g(z)|$ for all $z \in \mathbb{C}$. What conclusion can you draw?