


Colored
vertex models
and k-tilings of the
Aztec diamond

Sylvie Corteel (CNRS, IRIF)
UC Berkeley (on leave)




Calico - April 2022


C. , Gitlin , Keating & Meza
IMRN(2021)

Gitlin , Keating
Arxiv 2110.10273


C. , Gitlin & Keating
Arxiv 2202.06020



D. Keating
→ Wisconsin



J. Meza
→ Goldman Sachs




A. Gitlin
Berkeley


C. , Gittlin , Keating & Meza
IMRN(2021)

Gittlin , Keating
Arxiv 2110.10273


C. , Gittlin & Keating
Arxiv 2202.06020



D. Keating
→ Wisconsin



J. Meza
→ Goldman
Sachs



A. Gittlin
Berkeley

Many other papers
"Solvable lattice model seminar"
Tuesday 2pm PST

Five vertex models

(Folklore)


Partition of an integer n

$$\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$$

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k \geq 0$$

$$\lambda_1 + \dots + \lambda_k = n$$



Maya diagram of a partition




$$\lambda = (4, 3, 1, 0, 0)$$

... 0 0 0 0 0 0 0 0 0 0 ...


Five vertex model



Five vertex model



λ




μ

Interlacing $\lambda \succcurlyeq \mu$

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \lambda_3 \geq \mu_3 \geq \dots$$

Five vertex model



Interlacing $\lambda \succcurlyeq \mu$



$$\lambda_1 \succcurlyeq \mu_1 \succcurlyeq \lambda_2 \succcurlyeq \mu_2 \succcurlyeq \lambda_3 \succcurlyeq \mu_3 \succcurlyeq \dots$$

ex $\lambda = (5, 5, 2, 1)$ $\mu = (5, 3, 1, 0)$

$$5 \succcurlyeq 5 \succcurlyeq 5 \succcurlyeq 3 \succcurlyeq 2 \succcurlyeq 1 \succcurlyeq 1 \succcurlyeq 0$$

Weight x^4

Weight of a configuration




$$\text{wt}(c) = \prod_{\text{vertex}} \text{weight of the vertex}$$


ex

$$x_1^3 x_2^2 x_3 x_4$$

Partition function



$$Z_{\lambda/\mu}(x_1, x_2, \dots, x_n) = \sum_{\substack{c \text{ configurations} \\ \text{from } \mu \text{ to } \lambda}} \text{wt}(c)$$



$$\mu = (2, 2, 1, 1, 1)$$

$$\lambda = (4, 4, 4, 4, 2)$$

...

↖

		1	1
		2	2
	1	3	3
	2	4	4
	3		

↗

Prop

$$Z_{\lambda/\mu}(x_1, x_2, \dots, x_n) = S_{\lambda/\mu}(x_1, \dots, x_n)$$

Skew - Schur polynomials

Prop For any permutation $\pi \in S_n$

$$S_{\lambda/\mu}(x_1, \dots, x_n) = S_{\lambda/\mu}(x_{\pi_1}, \dots, x_{\pi_n})$$

Proof Yang Baxter equation


$$\sum_{\text{conf}} \begin{array}{c} \text{F} \\ \begin{array}{|c|} \hline x \\ \hline y \\ \hline \end{array} \\ \begin{array}{c} a \quad e \\ b \quad d \\ c \end{array} \end{array} = \sum_{\text{conf}} \begin{array}{c} \text{F} \\ \begin{array}{|c|} \hline y \\ \hline x \\ \hline \end{array} \\ \begin{array}{c} a \quad e \\ b \quad d \\ c \end{array} \end{array}$$

$$\forall a, b, c, d, e, F \in \{0, 1\}$$





Prop For any permutation $\pi \in S_n$



$$S_{\lambda/\mu}(x_1, \dots, x_n) = S_{\lambda/\mu}(x_{\pi_1}, \dots, x_{\pi_n})$$

Proof Yang Baxter equation





R






$$\underline{\lambda \succ \mu}$$




Pyramid partition (Young) Rail yard Graphs (BBCCR)



Cauchy identity.

$$\sum_{\lambda} s_{\lambda/\mu}(x) s_{\lambda/\nu}(y) = \frac{1}{1-xy} \sum_{\lambda} s_{\mu/\lambda}(y) s_{\nu/\lambda}(x)$$





Cauchy identity.

$$\sum_{\lambda} s_{\lambda/\mu}(x) s_{\lambda/\nu}(y) = \frac{1}{1-xy} \sum_{\lambda} s_{\mu/\lambda}(y) s_{\nu/\lambda}(x)$$




Dual Cauchy identity

$$\sum_{\lambda} s_{\lambda/\mu}(x) s_{(\lambda/\nu)'}(y) = (1+xy) \sum_{\lambda} s_{\mu/\lambda'}(y) s_{\nu/\lambda}(x)$$



Cauchy identity.


$$\sum_{\lambda} S_{\lambda/\mu}(x) S_{\lambda/\nu}(y) = \frac{1}{1-xy} \sum_{\lambda} s_{\mu/\lambda}(y) s_{\nu/\lambda}(x)$$



Dual Cauchy identity



$$\sum_{\lambda} S_{\lambda/\mu}(x) S_{(\lambda/\nu)'}(y) = (1+xy) \sum_{\lambda} s_{(\mu/\lambda)'}(y) s_{\nu/\lambda}(x)$$

Domino tilings of the Aztec diamond (Propp et al 90s)




m corners

Domino tilings of the Aztec diamond (Propp et al 90s)





Domino tilings of the Aztec diamond

$$\underline{m=2}$$




Theorem The number of domino Tilings of
the Aztec diamond with m corners
is $2^{\binom{m+1}{2}}$


Domino tilings of the Aztec diamond



Domino tilings of the Aztec diamond




Domino tilings of the Aztec diamond



$$\emptyset = \lambda^{(0)} \prec \lambda^{(1)} \succ \lambda^{(2)} \\ \dots \prec \lambda^{(2m-1)} \succ \lambda^{(2m)} \\ = \emptyset$$



$$Z_n(x, y) = \sum_{\lambda} \prod_i x_i^{\frac{|\lambda^{(2i-1)}|}{|\lambda^{(2i-2)}|}} y_i^{\frac{|\lambda^{(2i-1)}|}{|\lambda^{(2i)}|}} \\ = \prod_{1 \leq i \leq j \leq m} (1 + x_i y_j)$$


Aztec diamond $\rightarrow$ Vertex models



$$\prod_{i \leq j} (1 + x_i y_j)$$


Steep tilings
(Bouttier, Chapuy, C.)
2013










$\gamma B E$







=




weight₁
(1 + $x_i y_m$)





$=$




$$\times \frac{1}{\prod_{1 \leq i \leq j \leq m} (1 + x_i y_j)}$$


Colored five vertex models



(Five)² vertex model




(Five)² vertex model




(Five)^k relax models

Weight for color i


$i = 1, \dots, k$




1




1



1



$x t^{\otimes}$




$x t^{\otimes}$

$\otimes$: # larger colors that are present


Weight of a vertex = $\prod_{i=1}^k$ weight of color i

(Five)^k relax models


Weight for color i




1




1



1




$x t^{\oplus}$




$x t^{\oplus}$

$\oplus$: # larger colors that are present

$R < B < G$




Weight $x^2 t^2$



$x^2 t^{2+1}$

Partition function



$$Z_{\lambda/\mu}(x_1, x_2, \dots, x_n) = \sum_{\substack{c \text{ configurations} \\ \text{from } \mu \text{ to } \lambda}} \text{wt}(c)$$

Theorem (C., Gitlin, Keating & Meza 2020)

$Z_{\lambda, \mu}(x_1, \dots, x_n; t)$ is equal to the

L LT polynomial $Z_{\lambda, \mu}(x_1, \dots, x_n; t)$

$$\lambda = (\lambda^{(1)}, \dots, \lambda^{(k)})$$

$$\mu = (\mu^{(1)}, \dots, \mu^{(k)})$$

Theorem (C., Gitlin, Keating & Meza)


$Z_{\lambda, \mu}(x_1, \dots, x_n; t)$ is equal to the

LIT polynomial $Z_{\lambda/\mu}(x_1, \dots, x_n; t)$


$$\lambda = (\lambda^{(1)}, \dots, \lambda^{(k)})$$

$$\mu = (\mu^{(1)}, \dots, \mu^{(k)})$$


LIT polynomials (Haglund, Haiman, Loehr)






$$Z_{\lambda/\mu} = \sum_{T \text{ sequence of SSYT}} x^T t^{\text{coinv}(T)}$$




coinversion
 $u \leq v \leq w$











$$B < R$$









Weight $x_1^3 x_2^3 x_3^2 t^3$

Yang Baxter equation




a, b, c, d, e, f
fixed $\in \{0, 1\}^k$




♥ : # larger colors 

Yang Baxter equation



a, b, c, d, e, F
fixed $\in \{0, 1\}^k$



Proofs

Combinatorics (C., Gritsin, Keating, Vera)
Representation theory / Fusion
(Aggarwal, Bondin, Wheeler)

Corollaries of the Yang Baxter equation


① $L_{\lambda/\mu}(x_1, \dots, x_n; t)$ is a symmetric polynomial

② Cauchy identity (Generalization of Lam OS, Aggarwal 21)
C., GKP 2020 Borodin Wheeler

$$\prod_{i=0}^{k-1} (1 - xyt^i) \sum_{\lambda} t^{d(\lambda)} L_{\lambda/\mu}(x) L_{\lambda/\nu}(y) \\ = \sum_{\lambda} t^{d(\lambda)} L_{\mu/\lambda}(y) L_{\nu/\lambda}(x)$$


Generalisation to the base $(\text{five})^k$ vertex models

Supersymmetric LLT polynomials
(Gitten & Keating 2021)




Generalisation to the base (five)^k vertex models

Supersymmetric LLT polynomials
(Gitten & Keating 2021)




k-tilings of the Aztec diamond


$k=3$



T_1




T_2




T_3

k-tilings of the Aztec diamond


$k=3$




T_1



T_2




T_3



,



,




, or



.


k-tilings of the Aztec diamond

$k=3$




k-tilings of the Aztec diamond


$k=3$




T_1



T_2




T_3



,



,





, or



.


k-tilings of the Aztec diamond

$k=3$



k-tilings of the Aztec diamond

$k=3$



11 interactions

$$Z_m^{(k)}(x, y, t) = \sum_{\substack{k\text{-filings} \\ \text{of the} \\ m\text{-diamond}}} t^{\# \text{inclusions}} \prod_{i=1}^k \text{weight}^{(i)}(xy)$$

$$Z_m^{(k)}(x, y, t) = \sum_{k\text{-tilings}} t^{\# \text{ inversions}} \prod_{i=1}^k \text{weight}^{(i)}(xy)$$





Theorem (C., Ghitu, Keating 2022)



$$Z_m^{(k)}(x, y, t) = \prod_{\ell=0}^{k-1} \prod_{1 \leq i \leq j \leq m} (1 + t^{\ell} x_i y_j)$$

$$Z_m^{(k)}(x, y, t) = \sum_{k\text{-tuples}} t^{\# \text{inclusions}} \prod_{i=1}^k \text{weight}^{(i)}(xy)$$



Theorem (C., Githu, Keehye 2022)

$$Z_m^{(k)}(x, y, t) = \prod_{\ell=0}^{k-1} \prod_{i \leq j} (1 + t^{\ell} x_i y_j)$$

 T_1  T_2  T_3



Same partition function



Same partition function



$$\underline{t = 0}$$


$$Z_m^{(k)}(x, y, 0) = \prod_{1 \leq i \leq j \leq m} (1 + x_i y_j)$$

↑
1-tilings

$$\underline{t = 0}$$




$$Z_m^{(k)}(x, y, 0) = \prod_{i \leq j} (1 + x_i y_j) \\ = Z_m(x, y)$$

Tilings $\rightarrow$ Paths





$$\underline{t = 0}$$

$$Z_m^{(k)}(x, y, 0) = \prod_{i \leq j} (1 + x_i y_j) \\ = Z_m(x, y)$$



$$\underline{t = 0}$$

$$Z_m^{(k)}(x, y, 0) = \prod_{i \leq j} (1 + x_i y_j) \\ = Z_m(x, y)$$



Questions

- ⑥ k - Pyramid partitions?
- ① RSK?
- ② Domino shuffling?
(Keating & Nicodetti)
- ③ Kuo condensation?
- ④ Asymptotics $t \neq 0, 1, \infty$
Limit shape? Height function?

